

Oxalis stricta

Habitus and growth type

Plant height [m]: **0.22**

Life span: **Annual, Biennial or short-lived, Perennial**

Life form: **Hemicryptophyte, Geophyte, Therophyte**

Leaf

Specific leaf area [mm²/mg]: **57.7**

Flower

Flowering period: **April-November**

Fruit, seed and dispersal

Seed mass [mg]: **0.18**

Dispersal mode: **Local non-specific dispersal, Anthropochory**

Dispersal distance class: **1, 7**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **neophyte**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**

Substrate reaction relationship: **Slightly acidic to near-neutral**

Nutrient relationship: **Eutrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **6.5**

Temperature indicator value: **6.2**

Moisture indicator value: **5**

Reaction indicator value: **5.5**

Nutrient indicator value: **6.5**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **1.14**

Disturbance frequency (herb layer): **1.84**

Disturbance severity: **0.81**

Disturbance severity (herb layer): **0.77**

Mowing frequency: **0.4**

Grazing pressure: **0.14**

Soil disturbance: **0.7**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [QA \(PAR\) *Papaveretea rhoeadis*](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [V12 Mixed crops of market gardens and horticulture](#)

Constant species of EUNIS habitats: [V12 Mixed crops of market gardens and horticulture](#)

Broad habitat

Occurrence in broad habitats: **Sparsely vegetated (incl. rock and scree),**

Synanthropic

Distribution

Continentality amplitude: **5**