

# *Oxytropis campestris*

## Habitus and growth type

Plant height [m]: **0.21**

Life span: **Perennial**

Life form: **Hemicryptophyte**

## Leaf

Specific leaf area [mm<sup>2</sup>/mg]: **14.93**

## Flower

Flowering period: **May-August**

## Fruit, seed and dispersal

Seed mass [mg]: **2.14**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **1**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **symbiosis with rhizobia**

## Taxon origin

Origin in Europe: **Native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Dry**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **9**

Temperature indicator value: **2.3**

Moisture indicator value: **3.8**

Nutrient indicator value: **2.3**

Salinity indicator value: **0**

### Disturbance indicator values

Disturbance frequency: **1.52**

Disturbance frequency (herb layer): **1.65**

Disturbance severity: **0.33**

Disturbance severity (herb layer): **0.32**

Mowing frequency: **0.06**

Grazing pressure: **0.27**

Soil disturbance: **0.12**

## **Habitat and sociology**

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Alpine-subalpine and arctic grassland, Scrub, Sparsely vegetated (incl. rock and scree)**

## **Distribution**

Continentality: 7

Continentality amplitude: 5