

Parapholis marginata

Habitus and growth type

Plant height [m]: **0.13**

Life span: **Annual**

Life form: **Therophyte**

Flower

Flowering period: **March-June**

Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **1**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Eutrophic**

Ellenberg-type indicator values

Light indicator value: **8.5**

Temperature indicator value: **9.5**

Moisture indicator value: **5**

Reaction indicator value: **8**

Nutrient indicator value: **6**

Salinity indicator value: **6**

Disturbance indicator values

Disturbance frequency: **0.99**

Disturbance frequency (herb layer): **1.03**

Disturbance severity: **0.51**

Disturbance severity (herb layer): **0.46**

Mowing frequency: **0.13**

Grazing pressure: **0.19**

Soil disturbance: **0.31**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [JA \(SAG\) *Saginetea maritimae*](#)

Broad habitat

Occurrence in broad habitats: **Coastal saltmarsh, Coastal beach, dune or shingle, Saline vegetation, Sparsely vegetated (incl. rock and scree)**