

# *Parietaria cretica*

## Habitus and growth type

Plant height [m]: **0.18**

Life span: **Annual**

Life form: **Therophyte**

## Flower

Flowering period: **January-March**

## Fruit, seed and dispersal

Dispersal mode: **Epizoochory**

Dispersal distance class: **6**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **Native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Slightly acidic to near-neutral**

Nutrient relationship: **Eutrophic**

Salinity relationship: **Slightly saline or brackish**

### Ellenberg-type indicator values

Light indicator value: **7**

Temperature indicator value: **10.5**

Moisture indicator value: **4**

Reaction indicator value: **7.5**

Nutrient indicator value: **6.5**

Salinity indicator value: **0.5**

### Disturbance indicator values

Disturbance frequency: **0.72**

Disturbance frequency (herb layer): **1.33**

Disturbance severity: **0.7**

Disturbance severity (herb layer): **0.51**

Mowing frequency: **0.21**

Grazing pressure: **0.28**

Soil disturbance: **0.28**

## **Habitat and sociology**

Syntaxon

Diagnostic species of phytosociological classes: [JB \(CRI\) \*Crithmo-Staticetea\*](#), [KC \(ASP\) \*Asplenietea trichomanis\*](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [T25 \*Phoenix theophrasti\* vegetation](#)

Broad habitat

Occurrence in broad habitats: **Coastal cliff, Forest, Sparsely vegetated (incl. rock and scree)**