Parietaria judaica

Habitus and growth type

Plant height [m]: **0.27**
Life span: **Perennial**
Life form: **Hemicryptophyte**

Leaf

Specific leaf area [mm^2/mg]: **33.17**

Flower

Flowering period: **January-December**

Fruit, seed and dispersal

Seed mass [mg]: **0.23**
Dispersal mode: **Epizoochory**
Dispersal distance class: **6**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate reaction relationship: **Alkaline**
Nutrient relationship: **Eutrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **6.8**
Reaction indicator value: **7.6**
Nutrient indicator value: **6.3**
Salinity indicator value: **0.5**

Disturbance indicator values

Disturbance frequency: **0.79**
Disturbance frequency (herb layer): **1.3**
Disturbance severity: **0.7**
Disturbance severity (herb layer): **0.61**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Mowing frequency: **0.37**
Grazing pressure: **0.2**
Soil disturbance: **0.32**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **KD (CYM) *Cymbalario-Parietarietea diffusae***, **QC (CHE) *Chenopodietea***

EUNIS habitat

Diagnostic species of EUNIS habitats: **U38 Mediterranean base-rich inland cliff**, **U3D Wet inland cliff**

Constant species of EUNIS habitats: **T14 Mediterranean and Macaronesian riparian forest**, **U38 Mediterranean base-rich inland cliff**, **U3D Wet inland cliff**

Broad habitat

Occurrence in broad habitats: **Scrub, Forest, Sparsely vegetated (incl. rock and scree), Synanthropic**

Distribution

Continentality amplitude: **5**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz