

Parietaria judaica

Habitus and growth type

Plant height [m]: **0.27**

Life span: **Perennial**

Life form: **Hemicryptophyte**

Leaf

Specific leaf area [mm²/mg]: **33.17**

Flower

Flowering period: **January-December**

Fruit, seed and dispersal

Seed mass [mg]: **0.23**

Dispersal mode: **Epizoochory**

Dispersal distance class: **6**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Eutrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **6.8**

Reaction indicator value: **7.6**

Nutrient indicator value: **6.3**

Salinity indicator value: **0.5**

Disturbance indicator values

Disturbance frequency: **0.79**

Disturbance frequency (herb layer): **1.3**

Disturbance severity: **0.7**

Disturbance severity (herb layer): **0.61**

Mowing frequency: **0.37**

Grazing pressure: **0.2**

Soil disturbance: **0.32**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [KD \(CYM\) Cymbalario-Parietarietea diffusae](#), [QC \(CHE\) Chenopodietea](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [U3D Wet inland cliff](#)

Constant species of EUNIS habitats: [T14 Mediterranean and Macaronesian riparian forest](#), [U38 Mediterranean base-rich inland cliff](#), [U3D Wet inland cliff](#)

Broad habitat

Occurrence in broad habitats: **Scrub, Forest, Sparsely vegetated (incl. rock and scree), Synanthropic**

Distribution

Continentality amplitude: **5**