Paronychia macrosepala

Habitus and growth type

Plant height [m]: **0.03**
Life span: **Perennial**
Life form: **Hemicryptophyte**

Flower

Flowering period: **March-June**

Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **1**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Ellenberg-type indicator values

Light indicator value: **8**
Temperature indicator value: **7.9**
Moisture indicator value: **3.7**
Reaction indicator value: **8**
Nutrient indicator value: **5**
Salinity indicator value: **0.2**

Disturbance indicator values

Disturbance frequency: **0.92**
Disturbance frequency (herb layer): **1.26**
Disturbance severity: **0.58**
Disturbance severity (herb layer): **0.44**
Mowing frequency: **0.11**
Grazing pressure: **0.29**
Soil disturbance: **0.28**

Habitat and sociology

Syntaxon

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Diagnostic species of phytosociological classes: **FB (ROS) *Ononido-Rosmarinetea*, FJ (TRA) *Stipo-Trachynietea distachyae***

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub, Sparsely vegetated (incl. rock and scree)**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz