Persicaria dubia

Habitus and growth type

Plant height [m]: **0.47**
Life span: **Annual**
Life form: **Therophyte**

Leaf

Specific leaf area [mm^2/mg]: **34.38**

Flower

Flowering period: **June-October**

Fruit, seed and dispersal

Seed mass [mg]: **1.39**
Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Wet**
Substrate reaction relationship: **Slightly acidic to near-neutral**
Nutrient relationship: **Eutrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **6.7**
Temperature indicator value: **5.9**
Moisture indicator value: **8.1**
Reaction indicator value: **6.2**
Nutrient indicator value: **7**
Salinity indicator value: **0.2**

Disturbance indicator values

Disturbance frequency: **0.94**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Disturbance frequency (herb layer): **1.57**
Disturbance severity: **0.63**
Disturbance severity (herb layer): **0.57**
Mowing frequency: **0.26**
Grazing pressure: **0.21**
Soil disturbance: **0.26**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **QI (BID) *Bidentetea***

EUNIS habitat

Diagnostic species of EUNIS habitats: **Q61 Periodically exposed shore with stable, eutrophic sediments with pioneer or ephemeral vegetation**
Constant species of EUNIS habitats: **Q61 Periodically exposed shore with stable, eutrophic sediments with pioneer or ephemeral vegetation**

Broad habitat

Occurrence in broad habitats: **Wetland**

Distribution

Continentality amplitude: **5**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz