

# *Petrorhagia nanteuillii*

## Habitus and growth type

Plant height [m]: **0.36**

Life span: **Annual**

Life form: **Therophyte**

## Flower

Flowering period: **April-September**

## Fruit, seed and dispersal

Seed mass [mg]: **0.28**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **Native**

## Ecology

Environmental relationships

Substrate humidity relationship: **Dry**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **8.5**

Temperature indicator value: **7**

Moisture indicator value: **2.5**

Salinity indicator value: **1**

Disturbance indicator values

Disturbance frequency: **1.29**

Disturbance frequency (herb layer): **2.02**

Disturbance severity: **0.59**

Disturbance severity (herb layer): **0.56**

Mowing frequency: **0.41**

Grazing pressure: **0.35**

Soil disturbance: **0.3**

## Habitat and sociology

### Syntaxon

Diagnostic species of phytosociological classes: [FI \(TUB\) \*Helianthemetea guttati\*](#)

### EUNIS habitat

Diagnostic species of EUNIS habitats: [R24 Iberian summer pasture \(vallicar\)](#)

### Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub**