

Peucedanum palustre

Habitus and growth type

Plant height [m]: **0.93**

Life span: **Biennial or short-lived**

Life form: **Hemicryptophyte**

Leaf

Specific leaf area [mm²/mg]: **20.09**

Flower

Flowering period: **June-August**

Fruit, seed and dispersal

Seed mass [mg]: **2.42**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Wet**

Substrate reaction relationship: **Slightly acidic to near-neutral**

Nutrient relationship: **Mesotrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **6.9**

Temperature indicator value: **5.4**

Moisture indicator value: **8.8**

Reaction indicator value: **5.6**

Nutrient indicator value: **4.3**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.61**

Disturbance frequency (herb layer): **1.33**

Disturbance severity: **0.42**

Disturbance severity (herb layer): **0.23**

Mowing frequency: **0.24**

Grazing pressure: **0.14**

Soil disturbance: **0.12**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [IA \(ALN\) *Alnetea glutinosae*](#), [OD \(PHR\) *Phragmito-Magnocaricetea*](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [T15 Broadleaved swamp forest on non-acid peat](#)

Constant species of EUNIS habitats: [Q42 Extremely rich moss-sedge fen](#), [Q44 Calcareous quaking mire](#), [S92 Salix fen scrub](#), [T15 Broadleaved swamp forest on non-acid peat](#)

Broad habitat

Occurrence in broad habitats: **Aquatic, Wetland, Mire, Grassland (non-alpine, non-saline), Scrub, Forest**

Distribution

Continentality: **5**

Continentality amplitude: **4**