

# *Phelipanche nana*

## Habitus and growth type

Plant height [m]: **0.17**

Life span: **Annual, Perennial**

Life form: **Geophyte, Therophyte**

## Flower

Flowering period: **March-July**

## Fruit, seed and dispersal

Dispersal mode: **Anemochory**

Dispersal distance class: **5**

## Trophic mode

Parasitism and mycoheterotrophy: **obligate root parasite**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **Native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Mesotrophic**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **7**

Moisture indicator value: **3.7**

Reaction indicator value: **7.3**

Salinity indicator value: **0.3**

### Disturbance indicator values

Disturbance frequency: **0.82**

Disturbance frequency (herb layer): **1.63**

Disturbance severity: **0.83**

Disturbance severity (herb layer): **0.74**

Mowing frequency: **0.43**

Grazing pressure: **0.24**

Soil disturbance: **0.44**

## Habitat and sociology

### Syntaxon

Diagnostic species of phytosociological classes: [DA \(FES\) Festuco-Brometea](#)

### Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub**