# *Phelipanche purpurea*

### Habitus and growth type

Plant height [m]: **0.3**
Life span: **Perennial**
Life form: **Geophyte**

### Flower

Flowering period: **May-August**

### Fruit, seed and dispersal

Seed mass [mg]: **0.01**
Dispersal mode: **Anemochory**
Dispersal distance class: **5**

### Trophic mode

Parasitism and mycoheterotrophy: **obligate root parasite**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

### Taxon origin

Origin in Europe: **Native**

### Ecology

#### Environmental relationships

Substrate humidity relationship: **Dry**
Substrate reaction relationship: **Alkaline**
Nutrient relationship: **Oligotrophic**
Salinity relationship: **Non-saline**

#### Ellenberg-type indicator values

Light indicator value: **7.7**
Temperature indicator value: **6.6**
Moisture indicator value: **3.7**
Reaction indicator value: **7.2**
Nutrient indicator value: **2.3**
Salinity indicator value: **0**

#### Disturbance indicator values

Disturbance frequency: **0.76**
Disturbance frequency (herb layer): **1.59**
Disturbance severity: **0.53**
Disturbance severity (herb layer): **0.42**

*© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz*

Mowing frequency: **0.34**
Grazing pressure: **0.28**
Soil disturbance: **0.21**

## Habitat and sociology

### Syntaxon

Diagnostic species of phytosociological classes: **DA (FES) *Festuco-Brometea***

### Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub**

## Distribution

Continentality: **6**
Continentality amplitude: **6**

*© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz*