

Phleum ambiguum

Habitus and growth type

Life span: **Perennial**

Life form: **Hemicryptophyte**

Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Ellenberg-type indicator values

Light indicator value: **8.7**

Temperature indicator value: **6.4**

Moisture indicator value: **2.7**

Reaction indicator value: **6.5**

Nutrient indicator value: **2.2**

Salinity indicator value: **0.3**

Disturbance indicator values

Disturbance frequency: **1.07**

Disturbance frequency (herb layer): **1.89**

Disturbance severity: **0.43**

Disturbance severity (herb layer): **0.37**

Mowing frequency: **0.17**

Grazing pressure: **0.36**

Soil disturbance: **0.14**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [DA \(FES\) Festuco-Brometea](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [R14 Perennial rocky grassland of the Italian Peninsula](#), [S74 Central Mediterranean mountain hedgehog-heath](#)

Constant species of EUNIS habitats: [R14 Perennial rocky grassland of the Italian Peninsula](#)

[Peninsula, S74 Central Mediterranean mountain hedgehog-heath](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline),
Oromediterranean scrub**