

Phleum crypsoides

Habitus and growth type

Life span: **Annual**

Life form: **Therophyte**

Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **1**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**

Salinity relationship: **Saline**

Ellenberg-type indicator values

Light indicator value: **9**

Temperature indicator value: **12**

Moisture indicator value: **2**

Reaction indicator value: **9**

Nutrient indicator value: **6**

Salinity indicator value: **8**

Disturbance indicator values

Disturbance frequency: **0.63**

Disturbance frequency (herb layer): **0.79**

Disturbance severity: **0.43**

Disturbance severity (herb layer): **0.37**

Mowing frequency: **0.07**

Grazing pressure: **0.18**

Soil disturbance: **0.17**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [**JA \(SAG\) *Saginetea maritimae***](#)

Broad habitat

Occurrence in broad habitats: **Saline vegetation**