

# *Phoenix theophrasti*

## Habitus and growth type

Life span: **Perennial**

Life form: **Phanerophyte, Tree**

## Fruit, seed and dispersal

Dispersal mode: **Endozoochory**

Dispersal distance class: **6**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **Native**

## Ecology

Environmental relationships

Substrate humidity relationship: **Wet**

Salinity relationship: **Slightly saline or brackish**

Ellenberg-type indicator values

Light indicator value: **7**

Temperature indicator value: **12**

Moisture indicator value: **8**

Reaction indicator value: **8**

Nutrient indicator value: **7**

Salinity indicator value: **2**

Disturbance indicator values

Disturbance frequency: **0.39**

Disturbance frequency (herb layer): **1.75**

Disturbance severity: **0.7**

Disturbance severity (herb layer): **0.29**

Mowing frequency: **0.03**

Grazing pressure: **0.3**

Soil disturbance: **0.16**

## Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **[FD \(NER\) Nerio-Tamaricetea](#)**

## EUNIS habitat

Diagnostic species of EUNIS habitats: [T25 Phoenix theophrasti vegetation](#)

Constant species of EUNIS habitats: [T25 Phoenix theophrasti vegetation](#)

Dominant species of EUNIS habitats: [T25 Phoenix theophrasti vegetation](#)

## Broad habitat

Occurrence in broad habitats: **Scrub, Forest**