Pholiurus pannonicus

Habitus and growth type

Life span: **Annual**
Life form: **Therophyte**

Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **1**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**
Substrate reaction relationship: **Alkaline**
Nutrient relationship: **Oligotrophic**
Salinity relationship: **Saline**

Ellenberg-type indicator values

Light indicator value: **8.8**
Temperature indicator value: **6.8**
Moisture indicator value: **4.8**
Reaction indicator value: **8**
Nutrient indicator value: **2**
Salinity indicator value: **8.5**

Disturbance indicator values

Disturbance frequency: **2.2**
Disturbance frequency (herb layer): **2.2**
Disturbance severity: **0.46**
Disturbance severity (herb layer): **0.46**
Mowing frequency: **0.22**
Grazing pressure: **0.48**
Soil disturbance: **0.24**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **DB (FEP) *Festuco-Puccinellietea***

EUNIS habitat

Diagnostic species of EUNIS habitats: **R62 Continental inland salt steppe**

Broad habitat

Occurrence in broad habitats: **Saline vegetation**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz