

# *Phyteuma michelii*

## Habitus and growth type

Plant height [m]: **0.34**

Life span: **Perennial**

Life form: **Hemicryptophyte**

## Flower

Flowering period: **June-August**

## Fruit, seed and dispersal

Seed mass [mg]: **0.12**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **Native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Mesic**

Substrate reaction relationship: **Acidic**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **8**

Temperature indicator value: **2.7**

Moisture indicator value: **5**

Reaction indicator value: **2.5**

Nutrient indicator value: **2.5**

Salinity indicator value: **0**

### Disturbance indicator values

Disturbance frequency: **0.6**

Disturbance frequency (herb layer): **1.51**

Disturbance severity: **0.43**

Disturbance severity (herb layer): **0.28**

Mowing frequency: **0.08**

Grazing pressure: **0.27**

Soil disturbance: **0.12**

## **Habitat and sociology**

### Syntaxon

Diagnostic species of phytosociological classes: [CT \(TRI\) \*Juncetea trifidi\*](#)

### Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub, Sparsely vegetated (incl. rock and scree)**