

Phyteuma sieberi

Habitus and growth type

Plant height [m]: **0.1**

Life span: **Perennial**

Life form: **Hemicryptophyte**

Flower

Flowering period: **June-August**

Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **1**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **8**

Temperature indicator value: **2.3**

Moisture indicator value: **3**

Reaction indicator value: **8**

Nutrient indicator value: **2**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **1.01**

Disturbance frequency (herb layer): **1.21**

Disturbance severity: **0.28**

Disturbance severity (herb layer): **0.26**

Mowing frequency: **0**

Grazing pressure: **0.22**

Soil disturbance: **0.08**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [CU \(SES\) *Elyno-Seslerietea*](#), [KC \(ASP\) *Asplenietea trichomanis*](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [U36 Temperate high-mountain base-rich inland cliff](#)

Broad habitat

Occurrence in broad habitats: **Coastal cliff, Grassland (non-alpine, non-saline), Sparsely vegetated (incl. rock and scree)**