

# ***Picris rhagadioloides***

## **Habitus and growth type**

Plant height [m]: **0.29**

Life span: **Annual**

Life form: **Therophyte**

## **Flower**

Flowering period: **May-August**

## **Fruit, seed and dispersal**

Dispersal mode: **Anemochory**

Dispersal distance class: **5**

## **Trophic mode**

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## **Taxon origin**

Origin in Europe: **Native**

## **Ecology**

### **Environmental relationships**

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Eutrophic**

Salinity relationship: **Non-saline**

### **Ellenberg-type indicator values**

Light indicator value: **8**

Temperature indicator value: **8.8**

Moisture indicator value: **4**

Reaction indicator value: **7.3**

Nutrient indicator value: **7.7**

Salinity indicator value: **0**

### **Disturbance indicator values**

Disturbance frequency: **0.95**

Disturbance frequency (herb layer): **1.85**

Disturbance severity: **0.78**

Disturbance severity (herb layer): **0.5**

Mowing frequency: **0.21**

Grazing pressure: **0.34**

Soil disturbance: **0.31**

## Habitat and sociology

### Syntaxon

Diagnostic species of phytosociological classes: [FI \(TUB\) \*Helianthemetea guttati\*](#)

### EUNIS habitat

Diagnostic species of EUNIS habitats: [S63 Eastern garrigue](#), [S72 Eastern Mediterranean spiny heath \(phrygana\)](#)

### Broad habitat

Occurrence in broad habitats: **Coastal cliff, Grassland (non-alpine, non-saline), Scrub, Sparsely vegetated (incl. rock and scree)**