

Pilosella cymosa

Habitus and growth type

Plant height [m]: **0.47**

Life span: **Perennial**

Life form: **Hemicryptophyte**

Leaf

Specific leaf area [mm²/mg]: **22.33**

Flower

Flowering period: **April-July**

Fruit, seed and dispersal

Seed mass [mg]: **0.12**

Dispersal mode: **Anemochory**

Dispersal distance class: **5**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7.3**

Temperature indicator value: **6**

Moisture indicator value: **3.6**

Reaction indicator value: **7.5**

Nutrient indicator value: **2.5**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.6**

Disturbance frequency (herb layer): **1.54**
Disturbance severity: **0.51**
Disturbance severity (herb layer): **0.35**
Mowing frequency: **0.27**
Grazing pressure: **0.27**
Soil disturbance: **0.16**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [DA \(FES\) Festuco-Brometea](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [R17 Heavy-metal dry grassland of the Balkans](#), [R1K Balkan and Anatolian oromediterranean dry grassland](#)

Constant species of EUNIS habitats: [R17 Heavy-metal dry grassland of the Balkans](#), [R1K Balkan and Anatolian oromediterranean dry grassland](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Forest, Sparsely vegetated (incl. rock and scree)**

Distribution

Continentality: **5**
Continentality amplitude: **4**