Pimpinella major

Habitus and growth type

Plant height [m]: **0.75**
Life span: **Perennial**
Life form: **Hemicryptophyte**

Leaf

Specific leaf area [mm^2/mg]: **13.38**

Flower

Flowering period: **May-August**

Fruit, seed and dispersal

Seed mass [mg]: **2.14**
Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**
Substrate reaction relationship: **Alkaline**
Nutrient relationship: **Eutrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7.1**
Moisture indicator value: **5.4**
Reaction indicator value: **7**
Nutrient indicator value: **6.1**
Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.75**
Disturbance frequency (herb layer): **1.7**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Disturbance severity: **0.53**
Disturbance severity (herb layer): **0.43**
Mowing frequency: **0.6**
Grazing pressure: **0.16**
Soil disturbance: **0.28**

Habitat and sociology

EUNIS habitat

Diagnostic species of EUNIS habitats: **R22 Low and medium altitude hay meadow**, **R23 Mountain hay meadow**
Constant species of EUNIS habitats: **R23 Mountain hay meadow**

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Alpine-subalpine and arctic grassland, Scrub, Forest, Sparsely vegetated (incl. rock and scree)**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz