

# *Pimpinella villosa*

## Habitus and growth type

Life span: **Perennial**

Life form: **Hemicryptophyte**

## Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **Native**

## Ecology

Ellenberg-type indicator values

Light indicator value: **9**

Temperature indicator value: **7**

Moisture indicator value: **1**

Reaction indicator value: **7**

Nutrient indicator value: **2**

Disturbance indicator values

Disturbance frequency: **0.43**

Disturbance frequency (herb layer): **1.56**

Disturbance severity: **0.52**

Disturbance severity (herb layer): **0.28**

Mowing frequency: **0.23**

Grazing pressure: **0.27**

Soil disturbance: **0.23**

## Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [FI \(TUB\) \*Helianthemetea guttati\*](#), [PA \(SCH\) \*Scheuchzerio palustris\*-\*Caricetea fuscae\*](#)

Broad habitat

Occurrence in broad habitats: **Mire, Grassland (non-alpine, non-saline)**