

# *Pinguicula alpina*

## Habitus and growth type

Plant height [m]: **0.06**

Life span: **Perennial**

Life form: **Hemicryptophyte**

## Leaf

Specific leaf area [mm<sup>2</sup>/mg]: **27.86**

## Flower

Flowering period: **May-July**

## Fruit, seed and dispersal

Seed mass [mg]: **7.41**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **1**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Wet**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **8.3**

Temperature indicator value: **3.1**

Moisture indicator value: **8**

Reaction indicator value: **7.8**

Nutrient indicator value: **2.4**

Salinity indicator value: **0**

### Disturbance indicator values

Disturbance frequency: **0.79**

Disturbance frequency (herb layer): **1.04**  
Disturbance severity: **0.29**  
Disturbance severity (herb layer): **0.23**  
Mowing frequency: **0.08**  
Grazing pressure: **0.18**  
Soil disturbance: **0.07**

## Habitat and sociology

### Syntaxon

Diagnostic species of phytosociological classes: [OA \(MON\) \*Montio-Cardaminetea\*](#), [PA \(SCH\) \*Scheuchzerio palustris-Caricetea fuscae\*](#)

### EUNIS habitat

Diagnostic species of EUNIS habitats: [Q45 Arctic-alpine rich fen](#)

Constant species of EUNIS habitats: [Q45 Arctic-alpine rich fen](#)

### Broad habitat

Occurrence in broad habitats: **Aquatic, Spring, Wetland, Mire, Grassland (non-alpine, non-saline), Scrub**

## Distribution

Continentality: 7

Continentality amplitude: 4