

Pistacia terebinthus

Habitus and growth type

Plant height [m]:

Life span: **Perennial**

Life form: **Phanerophyte, Tree, Shrub**

Spinescence: **not spinescent**

Leaf

Specific leaf area [mm²/mg]: **10.83**

Flower

Flowering period: **February-June**

Fruit, seed and dispersal

Seed mass [mg]: **41.5**

Dispersal mode: **Endozoochory**

Dispersal distance class: **6**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Mesotrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **8**

Temperature indicator value: **9**

Moisture indicator value: **3.3**

Reaction indicator value: **6.8**

Nutrient indicator value: **3.3**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.11**
Disturbance frequency (herb layer): **1.34**
Disturbance severity: **0.69**
Disturbance severity (herb layer): **0.17**
Mowing frequency: **0.02**
Grazing pressure: **0.26**
Soil disturbance: **0.11**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [CB \(PUB\) *Quercetea pubescentis*](#), [FA \(QUI\) *Quercetea ilicis*](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [T1A Mediterranean thermophilous deciduous forest](#)

Constant species of EUNIS habitats: [T1A Mediterranean thermophilous deciduous forest](#), [T21 Mediterranean evergreen Quercus forest](#), [T3A Mediterranean lowland to submontane Pinus forest](#), [T3D Mediterranean Cupressaceae forest](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub, Forest, Sparsely vegetated (incl. rock and scree)**