Pittosporum tobira

Habitus and growth type

Plant height [m]: **3.5**
Life span: **Perennial**
Life form: **Phanerophyte, Tree, Shrub**

Flower

Flowering period: **February-June**

Fruit, seed and dispersal

Seed mass [mg]:
Dispersal mode: **Endozoochory, Anthropochory**
Dispersal distance class: **6, 7**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **neophyte**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**
Substrate reaction relationship: **Slightly acidic to near-neutral**
Nutrient relationship: **Mesotrophic**
Salinity relationship: **Slightly saline or brackish**

Ellenberg-type indicator values

Light indicator value: **7**
Temperature indicator value: **8.5**
Moisture indicator value: **3**
Reaction indicator value: **5**
Nutrient indicator value: **5**
Salinity indicator value: **1**

Habitat and sociology

EUNIS habitat

Diagnostic species of EUNIS habitats: **T29 Broadleaved evergreen plantation of non site-native trees**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Broad habitat

Occurrence in broad habitats: **Scrub**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz