

# ***Poa bulbosa***

## **Habitus and growth type**

Plant height [m]: **0.21**

Life span: **Perennial**

Life form: **Hemicryptophyte, Geophyte**

## **Leaf**

Specific leaf area [mm<sup>2</sup>/mg]: **24.33**

## **Flower**

Flowering period: **March-July**

## **Fruit, seed and dispersal**

Seed mass [mg]: **0.65**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **1**

## **Trophic mode**

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## **Taxon origin**

Origin in Europe: **Native**

## **Ecology**

### **Environmental relationships**

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Slightly acidic to near-neutral**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Non-saline**

### **Ellenberg-type indicator values**

Light indicator value: **8.2**

Temperature indicator value: **6.8**

Moisture indicator value: **2.8**

Reaction indicator value: **5.4**

Nutrient indicator value: **2**

Salinity indicator value: **0.6**

### **Disturbance indicator values**

Disturbance frequency: **0.94**

Disturbance frequency (herb layer): **1.83**

Disturbance severity: **0.56**

Disturbance severity (herb layer): **0.49**

Mowing frequency: **0.31**

Grazing pressure: **0.36**

Soil disturbance: **0.29**

## Habitat and sociology

### EUNIS habitat

Diagnostic species of EUNIS habitats: [R1C Desert steppe](#), [R1D Mediterranean closely grazed dry grassland](#), [S34 Balkan-Anatolian submontane genistoid scrub](#), [S67 Aralo-Caspian semi-desert](#), [U2A Crimean base-rich scree](#)

Constant species of EUNIS habitats: [N17 Black Sea coastal dune grassland \(grey dune\)](#), [R11 Pannonian and Pontic sandy steppe](#), [R12 Cryptogam- and annual-dominated vegetation on siliceous rock outcrops](#), [R13 Cryptogam- and annual-dominated vegetation on calcareous and ultramafic rock outcrops](#), [R14 Perennial rocky grassland of the Italian Peninsula](#), [R15 Continental dry rocky steppic grassland and dwarf scrub on chalk outcrops](#), [R17 Heavy-metal dry grassland of the Balkans](#), [R1B Continental dry grassland \(true steppe\)](#), [R1C Desert steppe](#), [R1D Mediterranean closely grazed dry grassland](#), [R1J Cyrno-Sardean oromediterranean siliceous dry grassland](#), [R1N Open Iberian supramediterranean dry acid and neutral grassland](#), [R1R Mediterranean to Atlantic open, dry, acid and neutral grassland](#), [R24 Iberian summer pasture \(vallicar\)](#), [R62 Continental inland salt steppe](#), [S34 Balkan-Anatolian submontane genistoid scrub](#), [S63 Eastern garrigue](#), [S67 Aralo-Caspian semi-desert](#), [S68 Semi-desert sand dune with sparse scrub](#), [S72 Eastern Mediterranean spiny heath \(phrygana\)](#), [S74 Central Mediterranean mountain hedgehog-heath](#), [S75 Eastern Mediterranean mountain hedgehog-heath](#), [S94 Semi-desert riparian scrub](#), [T33 Mediterranean mountain Abies forest](#), [T38 Mediterranean montane Cedrus forest](#), [T3D Mediterranean Cupressaceae forest](#), [U2A Crimean base-rich scree](#)

Dominant species of EUNIS habitats: [R1D Mediterranean closely grazed dry grassland](#)

### Broad habitat

Occurrence in broad habitats: **Coastal beach, dune or shingle, Grassland (non-alpine, non-saline), Alpine-subalpine and arctic grassland, Saline vegetation, Semi-desert, Oromediterranean scrub, Scrub, Forest, Sparsely vegetated (incl. rock and scree)**

## Distribution

Continentality amplitude: **10**