

# *Potamogeton lucens*

## Habitus and growth type

Plant height [m]: **2.1**

Life span: **Perennial**

Life form: **Hydrophyte**

## Leaf

Specific leaf area [mm<sup>2</sup>/mg]: **66.51**

## Flower

Flowering period: **May-August**

## Fruit, seed and dispersal

Seed mass [mg]: **5.13**

Dispersal mode: **Hydrochory**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Water**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Eutrophic**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **6.1**

Moisture indicator value: **11.9**

Reaction indicator value: **6.6**

Nutrient indicator value: **7.1**

Salinity indicator value: **0.1**

### Disturbance indicator values

Disturbance frequency: **1.62**

Disturbance frequency (herb layer): **1.62**

Disturbance severity: **0.25**

Disturbance severity (herb layer): **0.25**

Mowing frequency: **0.03**

Grazing pressure: **0.01**

Soil disturbance: **0.02**

## **Habitat and sociology**

### Syntaxon

Diagnostic species of phytosociological classes: [NB \(POT\) \*Potamogetonetea\*](#)

### EUNIS habitat

Diagnostic species of EUNIS habitats: [P3d Fresh-water submerged vegetation](#), [P3e Fresh-water nymphaeid vegetation](#), [V14 Inundated or inundatable cropland, including rice fields](#)

Dominant species of EUNIS habitats: [P3d Fresh-water submerged vegetation](#)

### Broad habitat

Occurrence in broad habitats: **Aquatic, Synanthropic**

## **Distribution**

Continentality amplitude: **8**