Potamogeton natans

Habitus and growth type

Plant height [m]: **0.62**
Life span: **Perennial**
Life form: **Hydrophyte**

Leaf

Specific leaf area [mm^2/mg]: **27.52**

Flower

Flowering period: **March-July**

Fruit, seed and dispersal

Seed mass [mg]: **5.73**
Dispersal mode: **Hydrochory**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Water**
Substrate reaction relationship: **Alkaline**
Nutrient relationship: **Mesotrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **6.4**
Temperature indicator value: **4.5**
Moisture indicator value: **11.3**
Salinity indicator value: **0.2**

Disturbance indicator values

Disturbance frequency: **1.62**
Disturbance frequency (herb layer): **1.62**
Disturbance severity: **0.25**
Disturbance severity (herb layer): **0.25**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Mowing frequency: **0.05**
Grazing pressure: **0.02**
Soil disturbance: **0.03**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **NB (POT) *Potamogetonetea***

EUNIS habitat

Diagnostic species of EUNIS habitats: **P3c Fresh-water large pleustophyte vegetation, P3e Fresh-water nymphaeid vegetation**
Constant species of EUNIS habitats: **P3c Fresh-water large pleustophyte vegetation, P3e Fresh-water nymphaeid vegetation**
Dominant species of EUNIS habitats: **P3e Fresh-water nymphaeid vegetation**

Broad habitat

Occurrence in broad habitats: **Aquatic, Wetland**

Distribution

Continentality amplitude: **8**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz