

Potentilla astracanica

Habitus and growth type

Life span: **Perennial**

Life form: **Hemicryptophyte**

Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **1**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**

Ellenberg-type indicator values

Light indicator value: **8.3**

Temperature indicator value: **5**

Moisture indicator value: **5.1**

Disturbance indicator values

Disturbance frequency: **1.74**

Disturbance frequency (herb layer): **1.92**

Disturbance severity: **0.47**

Disturbance severity (herb layer): **0.46**

Mowing frequency: **0.52**

Grazing pressure: **0.4**

Soil disturbance: **0.2**

Habitat and sociology

EUNIS habitat

Diagnostic species of EUNIS habitats: [R17 Heavy-metal dry grassland of the Balkans](#)

Constant species of EUNIS habitats: [R17 Heavy-metal dry grassland of the Balkans](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline)**