

Potentilla micrantha

Habitus and growth type

Plant height [m]: **0.06**

Life span: **Perennial**

Life form: **Hemicryptophyte**

Flower

Flowering period: **January-May, December**

Fruit, seed and dispersal

Seed mass [mg]: **0.55**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **1**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Mesotrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **5.6**

Temperature indicator value: **6.9**

Moisture indicator value: **4.1**

Reaction indicator value: **7**

Nutrient indicator value: **3.7**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.02**

Disturbance frequency (herb layer): **1.3**

Disturbance severity: **0.69**

Disturbance severity (herb layer): **0.12**

Mowing frequency: **0.01**

Grazing pressure: **0.2**

Soil disturbance: **0.1**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [CB \(PUB\) Quercetea pubescentis](#), [CL \(GER\) Trifolio-Geranietea sanguinei](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [T1G Alnus cordata forest](#)

Constant species of EUNIS habitats: [T1G Alnus cordata forest](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub, Forest**

Distribution

Continentality: **5**

Continentality amplitude: **3**