Potentilla recta

Habitus and growth type

Plant height [m]: **0.44**
Life span: **Perennial**
Life form: **Hemicryptophyte**

Leaf

Specific leaf area [mm^2/mg]: **17.32**

Flower

Flowering period: **May-July**

Fruit, seed and dispersal

Seed mass [mg]: **0.24**
Dispersal mode: **Myrmecochory**
Dispersal distance class: **3**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**
Substrate reaction relationship: **Alkaline**
Nutrient relationship: **Oligotrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **8.5**
Temperature indicator value: **6.9**
Moisture indicator value: **3.4**
Reaction indicator value: **6.4**
Nutrient indicator value: **2.2**
Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **1.07**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Disturbance frequency (herb layer): **1.82**
Disturbance severity: **0.54**
Disturbance severity (herb layer): **0.49**
Mowing frequency: **0.57**
Grazing pressure: **0.32**
Soil disturbance: **0.21**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **QF (ART) *Artemisietea vulgaris***

EUNIS habitat

Diagnostic species of EUNIS habitats: **S34 Balkan-Anatolian submontane genistoid scrub**
Constant species of EUNIS habitats: **S34 Balkan-Anatolian submontane genistoid scrub**

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Oromediterranean scrub, Scrub, Synanthropic**

Distribution

Continentality: **5**
Continentality amplitude: **4**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz