

Pseudorlaya pumila

Habitus and growth type

Plant height [m]: **0.1**

Life span: **Annual**

Life form: **Therophyte**

Leaf

Specific leaf area [mm²/mg]: **6.81**

Flower

Flowering period: **March-May**

Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **1**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Slightly acidic to near-neutral**

Nutrient relationship: **Mesotrophic**

Salinity relationship: **Slightly saline or brackish**

Ellenberg-type indicator values

Light indicator value: **9**

Temperature indicator value: **10**

Moisture indicator value: **3.3**

Reaction indicator value: **7**

Nutrient indicator value: **4.7**

Salinity indicator value: **3**

Disturbance indicator values

Disturbance frequency: **1.48**

Disturbance frequency (herb layer): **1.95**

Disturbance severity: **0.44**

Disturbance severity (herb layer): **0.41**

Mowing frequency: **0.04**

Grazing pressure: **0.18**

Soil disturbance: **0.54**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [FI \(TUB\) *Helianthemetea guttati*](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [N16 Mediterranean and Macaronesian coastal dune grassland \(grey dune\)](#)

Constant species of EUNIS habitats: [N16 Mediterranean and Macaronesian coastal dune grassland \(grey dune\)](#)

Broad habitat

Occurrence in broad habitats: **Coastal beach, dune or shingle, Grassland (non-alpine, non-saline)**