

Puccinellia tenuifolia

Habitus and growth type

Life span: **Perennial**

Life form: **Hemicryptophyte**

Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Wet**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Eutrophic**

Salinity relationship: **Saline**

Ellenberg-type indicator values

Light indicator value: **8**

Temperature indicator value: **9**

Moisture indicator value: **8**

Reaction indicator value: **8**

Nutrient indicator value: **7**

Salinity indicator value: **8**

Disturbance indicator values

Disturbance frequency: **1.63**

Disturbance frequency (herb layer): **1.65**

Disturbance severity: **0.49**

Disturbance severity (herb layer): **0.47**

Mowing frequency: **0.02**

Grazing pressure: **0.31**

Soil disturbance: **0.25**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [MF \(JUN\) *Juncetea maritimi*](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [R61 Mediterranean inland salt steppe](#)

Broad habitat

Occurrence in broad habitats: **Coastal saltmarsh, Saline vegetation**