

# ***Ranunculus aconitifolius***

## **Habitus and growth type**

Plant height [m]: **0.4**

Life span: **Perennial**

Life form: **Hemicryptophyte**

## **Leaf**

Specific leaf area [mm<sup>2</sup>/mg]: **36.19**

## **Flower**

Flowering period: **April-July**

## **Fruit, seed and dispersal**

Seed mass [mg]: **3.56**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

## **Trophic mode**

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## **Taxon origin**

Origin in Europe: **Native**

## **Ecology**

### **Environmental relationships**

Substrate humidity relationship: **Wet**

Substrate reaction relationship: **Slightly acidic to near-neutral**

Nutrient relationship: **Eutrophic**

Salinity relationship: **Non-saline**

### **Ellenberg-type indicator values**

Light indicator value: **6.6**

Temperature indicator value: **3.7**

Moisture indicator value: **7.5**

Reaction indicator value: **5.2**

Salinity indicator value: **0**

### **Disturbance indicator values**

Disturbance frequency: **0.29**

Disturbance frequency (herb layer): **1.36**

Disturbance severity: **0.59**

Disturbance severity (herb layer): **0.27**

Mowing frequency: **0.13**

Grazing pressure: **0.18**

Soil disturbance: **0.11**

## **Habitat and sociology**

### Syntaxon

Diagnostic species of phytosociological classes: [CM \(MOL\) \*Molinio-Arrhenatheretea\*](#)

### EUNIS habitat

Diagnostic species of EUNIS habitats: [R56 Montane to subalpine moist or wet tall-herb and fern fringe](#)

### Broad habitat

Occurrence in broad habitats: **Spring, Grassland (non-alpine, non-saline), Alpine-subalpine and arctic grassland, Forest**

## **Distribution**

Continentality: **4**

Continentality amplitude: **2**