

Ranunculus arvensis

Habitus and growth type

Plant height [m]: **0.26**

Life span: **Annual**

Life form: **Therophyte**

Leaf

Specific leaf area [mm²/mg]: **27.4**

Flower

Flowering period: **March-June**

Fruit, seed and dispersal

Seed mass [mg]: **12.14**

Dispersal mode: **Local non-specific dispersal, Anthropochory**

Dispersal distance class: **1, 7**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Eutrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **6.9**

Moisture indicator value: **4.2**

Reaction indicator value: **7.7**

Nutrient indicator value: **6.1**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **1.82**

Disturbance frequency (herb layer): **2.05**

Disturbance severity: **0.84**

Disturbance severity (herb layer): **0.84**

Mowing frequency: **0.53**

Grazing pressure: **0.15**

Soil disturbance: **0.79**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [QA \(PAR\) *Papaveretea rhoeadis*](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [V13 Arable land with unmixed crops grown by low-intensity agricultural methods](#)

Constant species of EUNIS habitats: [V13 Arable land with unmixed crops grown by low-intensity agricultural methods](#)

Broad habitat

Occurrence in broad habitats: **Wetland, Synanthropic**

Distribution

Continentality: **4**

Continentality amplitude: **4**