

Ranunculus repens

Habitus and growth type

Plant height [m]: **0.2**

Life span: **Perennial**

Life form: **Hemicryptophyte**

Leaf

Specific leaf area [mm²/mg]: **24.63**

Flower

Flowering period: **February-August**

Fruit, seed and dispersal

Seed mass [mg]: **2.34**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **1**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Wet**

Substrate reaction relationship: **Slightly acidic to near-neutral**

Nutrient relationship: **Eutrophic**

Salinity relationship: **Slightly saline or brackish**

Ellenberg-type indicator values

Light indicator value: **6.3**

Temperature indicator value: **5**

Moisture indicator value: **7.3**

Reaction indicator value: **5.8**

Nutrient indicator value: **6.5**

Salinity indicator value: **0.7**

Disturbance indicator values

Disturbance frequency: **1.03**

Disturbance frequency (herb layer): **1.81**

Disturbance severity: **0.54**

Disturbance severity (herb layer): **0.49**

Mowing frequency: **0.61**

Grazing pressure: **0.33**

Soil disturbance: **0.27**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [**CM \(MOL\) Molinio-Arrhenatheretea**](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [**R21 Mesic permanent pasture of lowlands and mountains**](#), [**R35 Moist or wet mesotrophic to eutrophic hay meadow**](#), [**R36 Moist or wet mesotrophic to eutrophic pasture**](#)

Constant species of EUNIS habitats: [**N1H Atlantic and Baltic moist and wet dune slack**](#), [**P2N Spring**](#), [**Q52 Small-helophyte bed**](#), [**Q53 Tall-sedge bed**](#), [**Q61**](#)

[**Periodically exposed shore with stable, eutrophic sediments with pioneer or ephemeral vegetation**](#), [**R21 Mesic permanent pasture of lowlands and mountains**](#), [**R22 Low and medium altitude hay meadow**](#), [**R34 Submediterranean moist meadow**](#), [**R35 Moist or wet mesotrophic to eutrophic hay meadow**](#), [**R36 Moist or wet mesotrophic to eutrophic pasture**](#), [**R55 Lowland moist or wet tall-herb and fern fringe**](#), [**S38 Temperate forest clearing scrub**](#), [**S91 Temperate riparian scrub**](#), [**T11 Temperate Salix and Populus riparian forest**](#), [**T12 Alnus glutinosa-Alnus incana forest on riparian and mineral soils**](#), [**T13 Temperate hardwood riparian forest**](#), [**U71 Unvegetated or sparsely vegetated gravel bar in montane and alpine regions**](#), [**V11 Intensive unmixed crops**](#), [**V12 Mixed crops of market gardens and horticulture**](#), [**V13 Arable land with unmixed crops grown by low-intensity agricultural methods**](#), [**V15 Bare tilled, fallow or recently abandoned arable land**](#), [**V39 Mesic perennial anthropogenic herbaceous vegetation**](#)

Broad habitat

Occurrence in broad habitats: [**Aquatic**](#), [**Spring**](#), [**Wetland**](#), [**Grassland \(non-alpine, non-saline\)**](#), [**Saline vegetation**](#), [**Scrub**](#), [**Forest**](#), [**Sparingly vegetated \(incl. rock and scree\)**](#), [**Synanthropic**](#)

Distribution

Continentality amplitude: **7**