

# *Reynoutria sachalinensis*

## Habitus and growth type

Plant height [m]: **2.84**

Life span: **Perennial**

Life form: **Geophyte**

## Flower

Flowering period: **June-October**

## Fruit, seed and dispersal

Seed mass [mg]: **0.56**

Dispersal mode: **Local non-specific dispersal, Anthropochory**

Dispersal distance class: **2, 7**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **neophyte**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Wet**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Eutrophic**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **6.8**

Temperature indicator value: **6.8**

Moisture indicator value: **6.6**

Reaction indicator value: **6.6**

Nutrient indicator value: **7.6**

Salinity indicator value: **0**

### Disturbance indicator values

Disturbance frequency: **0.69**

Disturbance frequency (herb layer): **1.46**

Disturbance severity: **0.64**

Disturbance severity (herb layer): **0.52**

Mowing frequency: **0.53**

Grazing pressure: **0.25**

Soil disturbance: **0.21**

## **Habitat and sociology**

Syntaxon

Diagnostic species of phytosociological classes: [QG \(EPI\) \*Epilobietea angustifolii\*](#)

Broad habitat

Occurrence in broad habitats: **Wetland, Scrub, Forest, Synanthropic**

## **Distribution**

Continentality: **3**

Continentality amplitude: **3**