Rhus coriaria

Habitus and growth type

Plant height [m]: **2.75**
Life span: **Perennial**
Life form: **Phanerophyte, Shrub**

Leaf

Specific leaf area [mm^2/mg]: **10.55**

Flower

Flowering period: **April-August**

Fruit, seed and dispersal

Seed mass [mg]: **12.91**
Dispersal mode: **Endozoochory**
Dispersal distance class: **6**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**
Substrate reaction relationship: **Slightly acidic to near-neutral**
Nutrient relationship: **Mesotrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **8.3**
Temperature indicator value: **7.7**
Moisture indicator value: **4.4**
Reaction indicator value: **6.3**
Nutrient indicator value: **4.7**
Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.31**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Disturbance frequency (herb layer): **1.38**
Disturbance severity: **0.67**
Disturbance severity (herb layer): **0.34**
Mowing frequency: **0.14**
Grazing pressure: **0.25**
Soil disturbance: **0.18**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **CB (PUB) *Quercetea pubescentis*, KH (DRY) *Drypidetea spinosae***

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub, Forest, Sparsely vegetated (incl. rock and scree)**

Distribution

Continentality: **6**
Continentality amplitude: **5**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz