

Rorippa amphibia

Habitus and growth type

Plant height [m]: **0.83**

Life span: **Perennial**

Life form: **Hemicryptophyte**

Leaf

Specific leaf area [mm²/mg]: **30.16**

Flower

Flowering period: **April-July**

Fruit, seed and dispersal

Seed mass [mg]: **0.73**

Dispersal mode: **Epizoochory**

Dispersal distance class: **6**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Water**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Eutrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7.1**

Moisture indicator value: **9.7**

Reaction indicator value: **6.8**

Nutrient indicator value: **7.7**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **1.4**

Disturbance frequency (herb layer): **1.75**

Disturbance severity: **0.34**

Disturbance severity (herb layer): **0.33**

Mowing frequency: **0.24**

Grazing pressure: **0.16**

Soil disturbance: **0.12**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [OD \(PHR\) *Phragmito-Magnocaricetea*](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [Q52 Small-helophyte bed](#)

Constant species of EUNIS habitats: [Q52 Small-helophyte bed](#), [Q61 Periodically exposed shore with stable, eutrophic sediments with pioneer or ephemeral vegetation](#)

Broad habitat

Occurrence in broad habitats: **Wetland, Grassland (non-alpine, non-saline), Sparsely vegetated (incl. rock and scree)**

Distribution

Continentality: **6**

Continentality amplitude: **6**