

# *Rosa pouzinii*

## Habitus and growth type

Plant height [m]:

Life span: **Perennial**

Life form: **Phanerophyte, Shrub**

## Leaf

Specific leaf area [mm<sup>2</sup>/mg]: **14.7**

## Flower

Flowering period: **March-June**

## Fruit, seed and dispersal

Seed mass [mg]: **11.17**

Dispersal mode: **Endozoochory**

Dispersal distance class: **6**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **Native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Slightly acidic to near-neutral**

Nutrient relationship: **Mesotrophic**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **8**

Moisture indicator value: **2.7**

Reaction indicator value: **6.7**

Nutrient indicator value: **3**

Salinity indicator value: **0**

### Disturbance indicator values

Disturbance frequency: **0.2**

Disturbance frequency (herb layer): **1.2**

Disturbance severity: **0.64**

Disturbance severity (herb layer): **0.16**

Mowing frequency: **0**

Grazing pressure: **0.23**

Soil disturbance: **0.08**

## **Habitat and sociology**

### Syntaxon

Diagnostic species of phytosociological classes: [CD \(RHA\) Crataego-Prunetea](#)

### Broad habitat

Occurrence in broad habitats: **Scrub, Forest**