

# *Rudbeckia laciniata*

## Habitus and growth type

Plant height [m]: **1.45**

Life span: **Perennial**

Life form: **Hemicryptophyte, Geophyte**

## Flower

Flowering period: **June-September**

## Fruit, seed and dispersal

Seed mass [mg]: **2.61**

Dispersal mode: **Myrmecochory, Anthropochory**

Dispersal distance class: **3, 7**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **Neophyte**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Wet**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Eutrophic**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **7.3**

Temperature indicator value: **6.4**

Moisture indicator value: **7.6**

Reaction indicator value: **7**

Nutrient indicator value: **7**

Salinity indicator value: **0**

### Disturbance indicator values

Disturbance frequency: **0.64**

Disturbance frequency (herb layer): **1.41**

Disturbance severity: **0.58**

Disturbance severity (herb layer): **0.45**

Mowing frequency: **0.32**

Grazing pressure: **0.2**

Soil disturbance: **0.15**

## **Habitat and sociology**

Syntaxon

Diagnostic species of phytosociological classes: [QG \(EPI\) \*Epilobietea angustifolii\*](#)

Broad habitat

Occurrence in broad habitats: **Wetland, Scrub, Sparsely vegetated (incl. rock and scree), Synanthropic**

## **Distribution**

Continentality amplitude: **6**