

Rumex confertus

Habitus and growth type

Plant height [m]: **0.72**

Life span: **Perennial**

Life form: **Hemicryptophyte**

Flower

Flowering period: **June-August**

Fruit, seed and dispersal

Seed mass [mg]: **2.86**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate reaction relationship: **Slightly acidic to near-neutral**

Nutrient relationship: **Eutrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7.6**

Temperature indicator value: **4.8**

Reaction indicator value: **6**

Nutrient indicator value: **6.7**

Salinity indicator value: **0.3**

Disturbance indicator values

Disturbance frequency: **1.38**

Disturbance frequency (herb layer): **1.77**

Disturbance severity: **0.59**

Disturbance severity (herb layer): **0.57**

Mowing frequency: **0.88**

Grazing pressure: **0.18**

Soil disturbance: **0.27**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [CM \(MOL\) Molinio-Arrhenatheretea](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline)**