

Ruppia maritima

Habitus and growth type

Plant height [m]: **0.25**

Life span: **Perennial**

Life form: **Hydrophyte**

Leaf

Specific leaf area [mm²/mg]: **51.81**

Flower

Flowering period: **April-September**

Fruit, seed and dispersal

Seed mass [mg]: **1.22**

Dispersal mode: **Epizoochory**

Dispersal distance class: **6**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Water**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Eutrophic**

Ellenberg-type indicator values

Light indicator value: **7.8**

Moisture indicator value: **11**

Reaction indicator value: **7.8**

Nutrient indicator value: **7**

Salinity indicator value: **7.4**

Disturbance indicator values

Disturbance frequency: **1.74**

Disturbance frequency (herb layer): **1.74**

Disturbance severity: **0.34**

Disturbance severity (herb layer): **0.34**

Mowing frequency: **0.1**

Grazing pressure: **0.14**

Soil disturbance: **0.18**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [MC \(RUP\) *Ruppiaetea maritimae*](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [P3a Brackish-water vegetation](#)

Constant species of EUNIS habitats: [P3a Brackish-water vegetation](#)

Dominant species of EUNIS habitats: [P3a Brackish-water vegetation](#)

Broad habitat

Occurrence in broad habitats: **Marine, Coastal saltmarsh, Aquatic, Saline vegetation**

Distribution

Continentality: **8**

Continentality amplitude: **8**