

Sagina procumbens

Habitus and growth type

Plant height [m]: **0.07**

Life span: **Perennial**

Life form: **Hemicryptophyte**

Leaf

Specific leaf area [mm²/mg]: **19.25**

Flower

Flowering period: **March-July**

Fruit, seed and dispersal

Seed mass [mg]: **0.02**

Dispersal mode: **Local non-specific dispersal, Anthropochory**

Dispersal distance class: **1, 7**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**

Substrate reaction relationship: **Slightly acidic to near-neutral**

Nutrient relationship: **Eutrophic**

Salinity relationship: **Slightly saline or brackish**

Ellenberg-type indicator values

Light indicator value: **7.1**

Temperature indicator value: **4.9**

Reaction indicator value: **6**

Salinity indicator value: **1**

Disturbance indicator values

Disturbance frequency: **1.9**

Disturbance frequency (herb layer): **2.12**

Disturbance severity: **0.6**

Disturbance severity (herb layer): **0.59**

Mowing frequency: **0.65**

Grazing pressure: **0.41**

Soil disturbance: **0.37**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [QE \(POL\) *Polygono-Poetea annuae*](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [V35 Trampled mesophilous grassland with annuals](#)

Constant species of EUNIS habitats: [V35 Trampled mesophilous grassland with annuals](#)

Broad habitat

Occurrence in broad habitats: **Coastal saltmarsh, Grassland (non-alpine, non-saline), Synanthropic**

Distribution

Continentality amplitude: 5