

Salix breviserrata

Habitus and growth type

Plant height [m]: **0.21**

Life span: **Perennial**

Life form: **Chamaephyte**

Flower

Flowering period: **May-July**

Fruit, seed and dispersal

Seed mass [mg]: **0.15**

Dispersal mode: **Anemochory**

Dispersal distance class: **5**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Wet**

Substrate reaction relationship: **Slightly acidic to near-neutral**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7.5**

Temperature indicator value: **2.7**

Moisture indicator value: **7**

Reaction indicator value: **5.5**

Nutrient indicator value: **2.5**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **1.04**

Disturbance frequency (herb layer): **1.34**

Disturbance severity: **0.33**

Disturbance severity (herb layer): **0.25**

Mowing frequency: **0.09**

Grazing pressure: **0.24**

Soil disturbance: **0.12**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [CR \(VIR\) *Betulo carpaticae-Alnetea viridis*](#), [KE \(THL\) *Thlaspietea rotundifolii*](#)

Broad habitat

Occurrence in broad habitats: **Spring, Wetland, Grassland (non-alpine, non-saline), Scrub, Sparsely vegetated (incl. rock and scree)**