

Salix foetida

Habitus and growth type

Plant height [m]: **0.85**

Life span: **Perennial**

Life form: **Phanerophyte, Shrub**

Flower

Flowering period: **May-July**

Fruit, seed and dispersal

Dispersal mode: **Anemochory**

Dispersal distance class: **5**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Wet**

Substrate reaction relationship: **Acidic**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **8**

Temperature indicator value: **2.8**

Moisture indicator value: **7**

Reaction indicator value: **4**

Nutrient indicator value: **3.3**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **1.18**

Disturbance frequency (herb layer): **1.6**

Disturbance severity: **0.34**

Disturbance severity (herb layer): **0.25**

Mowing frequency: **0.12**

Grazing pressure: **0.17**

Soil disturbance: **0.09**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [CR \(VIR\) *Betulo carpaticae-Alnetea viridis*](#)

Broad habitat

Occurrence in broad habitats: **Scrub**

Distribution

Continentality: **6**

Continentality amplitude: **2**