

Salix hastata

Habitus and growth type

Plant height [m]: **0.98**

Life span: **Perennial**

Life form: **Phanerophyte, Shrub**

Leaf

Leaf type (woody plants only): **Broad deciduous or semi-deciduous**

Leaf phenology (woody plants only): **Winter deciduous**

Flower

Flowering period: **May-July**

Fruit, seed and dispersal

Seed mass [mg]: **0.14**

Dispersal mode: **Anemochory**

Dispersal distance class: **5**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Mesotrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7.4**

Temperature indicator value: **3**

Reaction indicator value: **6.8**

Nutrient indicator value: **3.4**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.74**

Disturbance frequency (herb layer): **1.21**

Disturbance severity: **0.35**

Disturbance severity (herb layer): **0.21**

Mowing frequency: **0.04**

Grazing pressure: **0.16**

Soil disturbance: **0.07**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [CR \(VIR\) *Betulo carpaticae-Alnetea viridis*](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [U35 Boreal and Arctic base-rich inland cliff](#)

Constant species of EUNIS habitats: [U35 Boreal and Arctic base-rich inland cliff](#)

Broad habitat

Occurrence in broad habitats: **Spring, Alpine-subalpine and arctic grassland, Scrub, Sparsely vegetated (incl. rock and scree)**

Distribution

Continentality: **6**

Continentality amplitude: **5**