

Salix salviifolia

Habitus and growth type

Life span: **Perennial**

Life form: **Phanerophyte, Shrub**

Fruit, seed and dispersal

Dispersal mode: **Anemochory**

Dispersal distance class: **5**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Ellenberg-type indicator values

Light indicator value: **9**

Temperature indicator value: **6**

Moisture indicator value: **12**

Reaction indicator value: **6**

Nutrient indicator value: **5**

Salinity indicator value: **0.1**

Disturbance indicator values

Disturbance frequency: **0.62**

Disturbance frequency (herb layer): **1.66**

Disturbance severity: **0.58**

Disturbance severity (herb layer): **0.38**

Mowing frequency: **0.01**

Grazing pressure: **0.24**

Soil disturbance: **0.15**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [FD \(NER\) Nerio-Tamaricetea](#), [HB \(PUR\) Salicetea purpureae](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [U72 Unvegetated or sparsely vegetated gravel](#)

[bar in the Mediterranean region](#)

Broad habitat

Occurrence in broad habitats: **Scrub, Sparsely vegetated (incl. rock and scree)**