

Salix starkeana

Habitus and growth type

Plant height [m]: **0.73**

Life span: **Perennial**

Life form: **Phanerophyte, Shrub**

Leaf

Leaf type (woody plants only): **Broad deciduous or semi-deciduous**

Leaf phenology (woody plants only): **Winter deciduous**

Specific leaf area [mm²/mg]: **12.89**

Fruit, seed and dispersal

Dispersal mode: **Anemochory**

Dispersal distance class: **5**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Wet**

Substrate reaction relationship: **Acidic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7**

Temperature indicator value: **4.3**

Moisture indicator value: **6.9**

Reaction indicator value: **4**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.56**

Disturbance frequency (herb layer): **1.51**

Disturbance severity: **0.57**

Disturbance severity (herb layer): **0.35**

Mowing frequency: **0.38**

Grazing pressure: **0.19**

Soil disturbance: **0.15**

Habitat and sociology

EUNIS habitat

Diagnostic species of EUNIS habitats: [T3H Larix light taiga](#)

Constant species of EUNIS habitats: [T3H Larix light taiga](#)

Broad habitat

Occurrence in broad habitats: **Heathland, Scrub, Forest**

Distribution

Continentality: **7**

Continentality amplitude: **4**