Salix waldsteiniana

Habitus and growth type

Plant height [m]: **0.75**
Life span: **Perennial**
Life form: **Phanerophyte, Shrub**

Flower

Flowering period: **May-July**

Fruit, seed and dispersal

Seed mass [mg]: **0.1**
Dispersal mode: **Anemochory**
Dispersal distance class: **5**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**
Substrate reaction relationship: **Alkaline**
Nutrient relationship: **Mesotrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7**
Temperature indicator value: **3**
Moisture indicator value: **6**
Reaction indicator value: **8**
Nutrient indicator value: **4.5**
Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.3**
Disturbance frequency (herb layer): **0.47**
Disturbance severity: **0.4**
Disturbance severity (herb layer): **0.16**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Mowing frequency: **0.01**
Grazing pressure: **0.15**
Soil disturbance: **0.04**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **CR (VIR) *Betulo carpaticae-Alnetea viridis***

EUNIS habitat

Diagnostic species of EUNIS habitats: **S25 Subalpine and subarctic deciduous scrub**, **S26 Subalpine Pinus mugo scrub**
Constant species of EUNIS habitats: **S25 Subalpine and subarctic deciduous scrub**
Dominant species of EUNIS habitats: **S25 Subalpine and subarctic deciduous scrub**

Broad habitat

Occurrence in broad habitats: **Scrub**

Distribution

Continentality: **6**
Continentality amplitude: **2**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz